

### ATAGENIX LABORATORIES

## Catalog Number:ATMA10315Mo Anti 2019-nCoV Spike glycoprotein S1 monoclonal antibody

#### **Product Details**

#### **Summary**

Product name Anti 2019-nCoV Spike glycoprotein S1 monoclonal antibody

**description**This antibody was produced from a hybridoma resulting from the fusion of a

mouse myeloma with B cells obtained from a mouse immunized with purified,

recombinant SARS-CoV-2(2019-nCoV) Spike glycoprotein.

Accession # P0DTC2

Alternative names Spike glycoprotein, S glycoprotein, E2, Peplomer protein, Spike protein S1, Spike

protein S2, Spike protein S2', S

Stability &Storage Use a manual defrost freezer and avoid repeated freeze thaw cycles.

Store at 2 to 8 °C for one week .

Store at -20 to -80 °C for twelve months from the date of receipt.

**Spcificity** Recognizes SARS-CoV-2 Spike glycoprotein

**Isotype** IgG 1

**Host** Mouse

**Clonality** Monoclonal

**Clone No.** 19-J-16

**Conjugation** Unconjugated

Species reactivity Severe acute respiratory syndrome coronavirus 2 (2019-nCoV) (SARS-CoV-2)

Tested applications WB

Immunogen Recombinant SARS-CoV-2 (2019-nCoV) Spike protein fragment 1[679-833]

#### **Background**

Mediates fusion of the virion and cellular membranes by acting as a class I viral fusion protein. Under the current model, the protein has at least three conformational states: pre-fusion native state, pre-hairpin intermediate state, and post-fusion hairpin state. During viral and target cell membrane fusion, the coiled coil regions (heptad repeats) assume a trimer-of-hairpins structure, positioning the fusion peptide in close proximity to the C-terminal region of the ectodomain. The formation of this structure appears to drive apposition and subsequent fusion of viral and target cell membranes.

#### Product performance

Form Liquid



### **ATAGENIX LABORATORIES**

# Catalog Number:ATMA10315Mo Anti 2019-nCoV Spike glycoprotein S1 monoclonal antibody

**Buffer** PBS, pH7.4, containing 0.05% proclin300, 50% glycerol.

Concentration 0.42mg/ml

MW 142kDa

**Application** 

Dilution Range WB:1:1000~4000

Note

For research use only .Not for use in clinical diagnostic procedures.